
Pathway for Accessing Treatments for Non-Hospitalised Adultsvii with Mild COVID-19 (Effective 10th July 2023)

Supporting Information available via GP Gateway



Notes:

- I. C&W residents testing positive Out of Area (OOA) should be assessed by Respiratory@Home for Paxlovid and prescription issued to the OOA Pharmacy (or equivalent) service.
- II. OOA patients testing positive in C&W will need to be referred to the appropriate CMDU or equivalent within their area. Temporary registration with a C&W GP will not give access to medical notes and so triage and prescription cannot happen.
- III. Clinicians requesting LFT tests for patients potentially eligible for COVID-19 treatments should clearly annotate the request with "clinical priority group".
- IV. GP Practice teams/NHS 111 are not expected to make the final decision of whether the treatment is right for a patient or prescribe any treatment, rather to refer patients that meet the policy criteria in broad terms.
- V. The final decision about eligibility rests with the Respiratory@Home clinician.
- VI. The pathway is based on available information and clinical opinion. It will be reviewed and updated, if required, as services develop, and national policy is updated.
- VII. Please note that this is an Adult only service/pathway. For patients under 18 years: please liaise directly with the relevant Paediatric team: Respiratory@Home are unable to advise on such cases.

Eligible for COVID-19 Treatments: Treatment Choice and CMDU Responsibilities

Ensure comprehensive assessment of clinical appropriateness to guide treatment choice with reference to:

NICE TA 878 Casirivimab plus imdevimab, nirmatrelvir plus ritonavir, sotrovimab and tocilizumab for treating COVID-19

Note: Please refer to the electronic versions at all times.

	1 st line		2 nd line			
	PAXLOVID (SmPC)		SOTROVIMAB (SmPC)			
Clinical Check Requirements	1. Confirm exclusions including: o Pregnancy		Paxlovid is contra-indicated or unsuitable Confirm exclusions			
•	o Severe hepatic or renal impairment ¹		3. Use in pregnancy: where expected benefit to mother justifies the			
	o <u>Assessing for Paxlovid treatment</u>		risk to the foetus; liaise with appropriate specialist			
	o Interactions with COVID-19 antiviral therapies		4. Treatment is delivered within 5 days of symptom onset			
	o <u>Liverpool Drug Interactions Checker</u>		5. Counsel			
	2. Commence treatment within 5 days of symptom onset		6. Issue PIL (<u>Sotrovimab</u>); <u>National patient website</u>			
	3. Counsel, specifically regarding:					
	o Pregnancy and contraception					
	o Gastro-intestinal side effects					
	o Completion of course, missed dose					
	4. Issue PIL (Paxlovid); UKHSA leaflet; National patient website					
	1. CMDUs to contact local services for pharmacy/specialist advice as needed					
	2. Prescribe treatment as appropriate:		I			
Advice	Paxlovid: Via NHS Community Pharmacies		Sotrovimab: Via Trust Pharmacy Departments. This will be provided			
	I. Discuss with patient:		from the CMDUs at George Eliot Hospital and Warwick hospital.			
	a. Preferred community pharmacy considering urgency of prescription.					
	b. Standard prescription charges are applicable.		Process TBC once known			
	c. Contacting the pharmacy to confirm when the prescription is ready.					
	d. Arranging collection by an asymptomatic patient representative.					
	II. Telephone the identified community pharmacy to:					
	a. Confirm ability to fulfil prescription before sending via EPS.					
	b. Confirm clinical checks have been completed.					
	c. Community Pharmacists are also responsible for a clinical check and may request additional					
	information. If requested, supply (via secure nhs.net email) the completed referral form. See page 3 for further details regarding contacting community pharmacies and arrangements for urgent					
	supply. d. Advise on patient plans for medication collection.					
	III. Complete electronic FP10 and send to selected community pharmacy:					
	a. Temporary EPS nomination may be necessary, or an EPS token provided					
	b. Include statement that it is a 'Covid Medicine Service Prescription' and telephone number for					
	queries in FP10 'pharmacy info' section.					
	3. Activity Reporting to ICB					
Notes	i. Where a decision to prescribe is reversed, please ensure that both the provider and community pharmacy, as appropriate, are notified.					
notes	ii. Combination treatment with an nMAB and an antiviral is NOT routinely recommended					
	iii. Patients who have previously received treatment with an nMAB or antiviral, & who meet the eligibility criteria, may receive treatment for a subsequent episode, if clinically appropriate					
	iv. Where supply availability is impacting choice of treatment, consider MDT discussion to agree optimum treatment for the patients' circumstances.					
e SmPC for dosing	modification for moderate renal impairment.					

Eligible for COVID-19 Treatments: Treatment Provision & Pharmacist Responsibilities

	PAXLOVID (SmPC)	SOTROVIMAB (SmPC)
Responsibilities	Via selected Community Pharmacies:	
and Actions	These pharmacies are requested to keep 2 packs of Paxlovid in stock.	Via Trust Pharmacy departments: Details TBC when known
	Respiratory@Home have been supplied with direct contact details of pharmacies.	
Treatment	1. Draw down prescription from the spine.	
provision	2. Confirm:	Further details to be added once confirmed
	a. prescription has been generated by the triaging clinician	
	b. clinical check has been completed by prescriber	
	c. there are no outstanding concerns to preclude use of Paxlovid	
	3. Pharmacy Clinical Check and Dispensing Pharmacists are responsible for a clinical check.	
	Additional clinical information including the completed Checklist may be requested from the	
	prescriber, who will supply via secure nhs.net email.	
	4. Provision of Paxlovid Patient Information Leaflet (PIL), additional UKHSA Paxlovid leaflet and a pulse	
	oximeter and further patient information (if the patient agrees with the triaging clinician to be	
	monitored by the Respiratory@Home team). (Community pharmacies can have oximeters supplied	
	by Respiratory@Home team).	
	5. Counsel patient in line with Paxlovid PIL, specifically regarding:	
	i. Pregnancy and contraception	
	ii. Gastro-intestinal side effects	
	iii. Completion of course, and missed dose	
	6. Collect standard prescription charge if applicable	

Resource Signposting

- In addition to resources on pages 1 and 2 of this document, a complete list of resources are available via <u>Liverpool COVID-19 Interactions (covid19-druginteractions.org)</u> including:
- Guidance for resuming paused or dose-adjusted comedications with Paxlovid
- Swallowing Difficulties Administering Paxlovid as Crushed Tablets
- Guidance for Paxlovid dosing in patients with renal disease and patients on dialysis
- COVID-19 Antivirals Community Pharmacy Coventry and Warwickshire
- 3. Specialist Pharmacy Service Medicines Advice contact details: telephone 0300 770 8564 or email: asksps.nhs@sps.direct
- 4. E-Learning training materials: COVID-19 Treatments is available here (Registration on NHS Learning Hub is required)
- 5. PAXLOVID ▼ (nirmatrelvir/ritonavir) PfizerPro UK
- 6. Key Opinion Leader videos including: managing drug-drug interactions, importance of managing COVID, identifying & educating high risk patients
- 7. Specialist Pharmacy Service: Accessing COVID-19 Medicines Advice for Primary Care

<u>COVID-19 Therapies – Dose Recommendations for Patients with Renal Impairment</u>

COVID-19 Therapies - Dose Recommendations for Patients with Hepatic Impairment